



## KEY BENEFITS

- Preventively limits growth and development of diseases in crops
- Activates plants own defense system
- Helps to fight against various plant pathogens
- Beneficial for agricultural crops
- Effective even in harsh weather conditions
- Safe for environment and farmers.


## USAGE GUIDELINES

### BROADCASTING



Mix the recommended quantity of bio- products in 50 kg well decomposed FYM/Compost and broadcast in the field before last inter-culturing operations before sowing.

### FERTIGATION



It is completely soluble in water, giving advantage for easy and labor-saving application through drip irrigation system. Mix recommended dose of formulation in ample amount of water and apply to the desired area.

### SPECIAL CARE SHOULD BE TAKEN WHEN TANK MIXING

- Use appropriate dosages of formulation as per recommendation.
- Partially fill the spray tank with clean water and then add the specific amount of Bio-product to the tank. Add the remaining water and mix THOROUGHLY.
- DO NOT LET MIXTURE STAND OVERNIGHT.
- DO NOT mix with copper-based fungicides, bactericides, concentrated acids such as sulfuric acid, solvents and oxidizing agents.
- Application in the early morning OR evening creates conditions favourable for germination of spores, giving better results.

### MICROBIAL CONSORTIA FOR HEALTHY GROWTH OF PLANTS

Best to include Rubitrol in IPM programme, when conditions are conducive to heavy disease pressure.

Preventively limits many foliar and soil borne fungal pathogens

**Formulation** : Powder (Soluble Free Flowing Powder)  
**Shelf life** : 2 years  
**Packing** : 500g & 1 Kg

### BROADCASTING



Mix the recommended quantity of bio- products in 50 kg well decomposed FYM/Compost and broadcast in the field before last inter-culturing operations before sowing.

### SEED TREATMENT



Can be applied as water-based slurry to cover seeds uniformly as per the weight of seeds.

### DRENCHING



Calculate the volume of water required for drenching to the number of plants in the desired area. Add the recommended dose of formulation in the water and drench it in the root zone.